


JOURNAL OF ENVIRONMENTAL MICROBIOLOGY AND TOXICOLOGY

Website: <http://journal.hibiscuspublisher.com/index.php/JEMAT/index>



MIC and NIC Values Determination of *Allium sativum* Extract Against *Aeromonas hydrophila* Using the Lambert-Pearson Model

Ahmad Syazwan Ismail¹, Mohd Yunus Shukor² and Noor Azlina Masdor^{1*}

¹Biotechnology and Nanotechnology Research Centre, Malaysian Agricultural Research and Development Institute (MARDI), Persiaran MARDI-UPM, 43400 Serdang, Selangor, Malaysia.

²Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, D.E, Malaysia.

*Corresponding author:

Dr. Noor Azlina Masdor,

Biotechnology and Nanotechnology Research Centre,
Malaysian Agricultural Research and Development Institute (MARDI),
Persiaran MARDI-UPM,
43400 Serdang,
Selangor,
Malaysia.

Email: azlina@mardi.gov.my

HISTORY

Received: 24th May 2022
Received in revised form: 4th July 2022
Accepted: 15th July 2022

KEYWORDS

MIC
NIC
Lambert-Pearson model
Allium sativum
Aeromonas hydrophila

ABSTRACT

Plant extracts have been studied extensively for their ability to suppress the growth of bacteria. Because these findings were not reported on in the benchmark values, it was not possible to compare the results of this study to other investigations. Nonlinear regression analysis of the data is one of the most accurate methods of determining this value. The Minimum Non-inhibitory concentration (NIC) and inhibitory concentration (MIC) values of the methanolic extract of *Allium sativum* against the pathogen *Aeromonas hydrophila* were successfully determined using Lambert and Pearson's modified Gompertz model. Using nonlinear regression is useful since the 95% confidence interval of the MIC and NIC can now be estimated. As shown by the model's high correlation coefficient (0.995), garlic's MIC and NIC values (28.02 mg/mL, 95% Confidence Interval from 26.42 to 29.71 mg/mL) and (12.60 mg/mL, 95% Confidence Interval from 11.88 to 13.37 mg/mL) suggest that it could be used as an inhibitory agent against this essential fish pathogen.

INTRODUCTION

Aeromonas hydrophila, a Gram-negative rod-shaped bacterium from the family Aeromonadaceae, is a bacilli-like bacteria. In addition to being exceedingly mobile, it can be found in soil, sewage, and brackish water. It contains a single polar flagellum. Adhesion, cytotoxins, lipases, and biofilm development are some of the virulence factors of bacteria, which involve their potential to produce a variety of simultaneous attacks on the bacterial system. Motile aeromonad septicaemia (MAS) has been linked to MAS in a number of freshwater fish species, and it is also thought to be spread through inadvertent scrapes [1]. Several countries throughout the world have documented cases of this virus, particularly affecting fish species, including the United States. Affected species include channel cat fish, hybrid striped bass, Tilapia (*Tilapia nilotica*), Snakehead fish (*Ophiocephallus striatus*), Goldfish (*Carassius auratus*), American eel (*Anguilla Rostrata*), Carp (*Cyprinus carpio*), Chinook salmon (*Oncorhynchus tshawytscha*) and Rainbow trout (*Oncorhynchus mykiss*), to name a few [2].

Several popular plant extracts, including *Allium sativum* or garlic, were evaluated against *A. hydrophila* in a recent study. In terms of medical value, garlic is the most frequently cited herb [2]. The use of garlic as a folk medicine has been documented in Ayurvedic and related Indian and Chinese ancient systems of medicine. There has been a resurgence in the use of herbal medicines in recent years, and eco-friendly phytoproducts are being researched as effective antifungal agents for human therapy.

The bacterium responds well to this plant extract's antimicrobial properties [2]. The determination of the IC₅₀ which was not determined in the publication but was later determined [3]. However, IC₅₀ values are one of several important parameters of efficacy of an antibiotic agent against pathogens. Two other important parameters are Minimum Non-inhibitory concentration (NIC) and inhibitory concentration (MIC), respectively. It is the lowest concentration of antibiotic (typically in g/mL) that inhibits growth in vitro. MIC An antimicrobial's ability to suppress bacterial growth increases with a lower

minimum inhibitory concentration (MIC). It is the smallest amount of an antibiotic that "fully" inhibits bacterial growth at the lowest concentration. Antibiotics that limit bacterial growth can be found in the NIC form. Growth is equal to the control at concentrations below the NIC. When defining "totally retarding bacterial growth" and "slowing bacteriological growth," it appears that the definitions were made on the fly [4].

A semi-quantitative test approach is used to estimate the lowest concentration of antimicrobial necessary to suppress bacteria growth in order to arrive at the MIC. To evaluate the preservative, microorganisms were inoculated into growth broth that included a small amount of preservative. Finally, the least concentration of antimicrobial that generated a clear solution, meaning no visible growth was found in this test [5,6]. Currently, microtiter plates are being used in place of the old-fashioned tubes. End-point indicators can be utilized when turbidity of a test substance interferes with a test. These include resazurin [4] and fluorescein diacetate [7]. Even though no increase was observed in one well, it was regarded as the MIC [8]. Numerous antibiotic investigations have been hindered by the lack of a quantifiable standard procedure. [9,10]. However, the principal problem encountered is that all MIC techniques currently used are semi-quantitative. Nonlinear regression was used by Lambert and Pearson to determine the NIC and MIC. The MIC and NIC are derived from the slope and inflection point using the modified Gompertz model above. [11]. Using nonlinear regression is useful since the 95% confidence interval of the MIC and NIC can now be estimated.

MATERIALS AND METHODS

Acquisition of Data

Data from the works of Ramena et al. [2], from figure 3 graphs were scanned and electronically processed using Webplotdigitizer 2.5 [12]. Using the software, data from scanned images is converted into a table with comma-separated columns. [13].

Measurement of NIC and MIC: Fitting of a modified Gompertz function

A modified Gompertz equation requires log concentration data and a y response that has been translated into a fractional unit, such as fractional area or another fraction of unity, in order to be used for data analysis (Eqn. 1).

$$y = A + Ce^{-3^{B(x-M)}} \quad (\text{Eqn. 1})$$

where A, B, C and M represents the y lower asymptote with a value of approximately zero, slope parameter, distance from the upper and lower asymptote (with a value of, approximately one) and log concentration of the inflexion point, respectively. The NIC and MIC (Eqns. 2 and 3) values are obtained through the intersection of the lines $y=A+C$ and $y=A$, with the equation of the line tangential to the point $(M, (A, A+Ce^{-1}))$, respectively [11].

$$MIC = 10^{(M+\frac{1}{B})} \quad (\text{Eqn. 2})$$

$$NIC = 10^{(M-\frac{1.718}{B})} \quad (\text{Eqn. 3})$$

RESULTS AND DISCUSSION

The antimicrobial's minimum inhibitory concentration (MIC) is one of these microbiological parameters. It has been widely accepted for a long time. There have been very few instances of this finding over the years, but it is now showing up more regularly in the results of routine tests. However, the ability to use it for effective and optimal therapy is still limited and sometimes, despite much higher costs incurred than in qualitative methods, it is completely unused. The calculated MIC and NIC values are shown in **Table 1** whilst the fitted curve to the inhibition data show good fitting ($R^2=0.995$) (**Fig. 1**). The MIC and NIC values.

Table 1. MIC and NIC values of methanolic extract of *A. sativum* against *A. hydrophila*.

	mg/mL	95% Confidence interval
MIC	28.02	26.42 to 29.71
NIC	12.60	11.88 to 13.37




Fig. 1. Fitting of the inhibitory effect of *A. sativum* against *A. hydrophila* using the Lambert-Pearson model.

Allium sativum is a potent antipathogen. In one study, the effect of the fresh extracts of *A. sativum* as an anticryptococcal agent in broth showed an MIC values between 125 and 250 $\mu\text{g/mL}$ [14]. *A. hydrophila* is a fish pathogen which can cause fish production to be affected. Numerous studies have been conducted to find potential inhibition agents from plants, animal extracts, synthetic chemicals and also nanomaterials. In one such study, the methanolic gray nail extract (*Nephrolepis biserrata*) shows an MIC value for *A. hydrophila* at 50 mg/mL [15]. Cinnamaldehyde, a plant-derived ingredient shows an MIC value against *A. hydrophila* of 256 $\mu\text{g/mL}$ [16]. Peppers are also known as a traditional bacterial growth inhibitory agent. In one study, five Brazilian Piper species (*Piper aduncum*, *Piper callosum*, *Piper hispidinervum*, *Piper hispidum* and *Piper marginatum*) exhibits MIC values ranging from 0.23 to 30 mg/mL [17]. Nanoparticles are also emerging antibiotics for pathogens. In one study, titanium dioxide (TiO_2) nanoparticle (NP) shows an MIC value against *Aeromonas hydrophila* at 20 $\mu\text{g/mL}$ [16] paving the way for the use of nanometals in aquaculture as antibiotics.

Antibiotics like oxytetracycline, sulfadimethoxine, and florfenicol, which are effective yet expensive, have prompted researchers to conduct exploratory searches for new antibiotics. H₂O₂, a chemical that is both safe for the environment and powerful against this bacterium, is one such option. However, prolonged usage of this chemical can harm aquatic organisms such as algae and zooplankton [18]. Plant-derived bioactive molecules can be used as an alternative to synthetic chemicals.[19,20]. Plant bioactive chemicals have been proven in numerous research to have antibacterial and antifungal activities. Thirty-one Brazilian plant methanolic extracts were found poisonous to *F. columnare* and *A. hydrophila* in one such study [21]. *Citrobacter freundii*, *Vibrio parahaemolyticus*, *Edwardsiella tarda*, *Staphylococcus aureus*, *Vibrio vulnificus*, *Escherichia coli*, and *Streptococcus agalactia* have been discovered to be inhibited by common plant extracts. [22]. The antibacterial activity of common spices like clove, garlic and dill is demonstrated even by the more popular ones like coriander, onion and cumin. [23] and is likely to be explored for their antibiotic properties against this important fish pathogen in the near future.

CONCLUSION

There is a lack of mathematical models or nonlinear regression used to obtain the MIC and NIC values, which are critical for comparison, efficacy, and validation studies, in many studies on plant extracts and bacterial pathogens. There is a lack of mathematical models or nonlinear regression used to obtain the MIC and NIC values, which are critical for comparison, efficacy, and validation studies, in many studies on plant extracts and bacterial pathogens. In this study, the modified Gompertz model introduced by Lambert and Pearson has been successfully used to determine the MIC and NIC values of the methanolic extract of *Allium sativum* against the pathogen *A. hydrophila*. The model gave a correlation coefficient value of 0.995, indicating acceptable fitting and MIC and NIC values of 28.02 and 12.60 mg/mL, respectively, indicating the potential utility of garlic as inhibitory agent for this important fish pathogen.

REFERENCES

- Hoque F. Biocontrol of β -haemolytic *Aeromonas hydrophila* infection in *Labeo rohita* using antagonistic bacterium *Pseudomonas aeruginosa* FARP72. *Int J Pharm Sci Rev Res*. 2014;5(2):490–501.
- Ramena G, Ramena Y, Challa N. Identification and determination of minimum inhibitory concentrations of plant extracts having antimicrobial activity as potential alternative therapeutics to treat *Aeromonas hydrophila* infections. *J Microb Pathog*. 2018 Jan 27;2(1):1–9.
- Rusnam. Determination of the IC₅₀ Value of *Allium sativum* methanolic extract against *Aeromonas hydrophila*. *J Biochem Microbiol Biotechnol*. 2018 Jul 31;6(1):19–21.
- Mann CM, Markham JL. A new method for determining the minimum inhibitory concentration of essential oils. *J Appl Microbiol*. 1998;84(4):538–44.
- Collins CH. Antibiotics and antibacterial substances. *Microbiol Methods*. 1964;296–305.
- Davidson PM, Parish ME. Methods for testing the efficacy of food antimicrobials. *Food Technol*. 1989;43(1):148–55.
- Chand S, Lusunzi I, Williams LR, Karuso P, Veal DA. Rapid screening of the antimicrobial activity of extracts and natural products. *J Antibiot (Tokyo)*. 1994;47(11):1295–304.
- Sommers HM. Drug susceptibility testing in vitro: Monitoring of antimicrobial therapy. *Biol Clin Basis Infect Dis*. 1980;782–804.
- Janssen AM, Scheffer JJC, Baerheim Svendsen A. Antimicrobial activity of essential oils: A 1976-1986 literature review. Aspects of the test methods. *Planta Med*. 1987;53(5):395–8.
- Manou I, Bouillard L, Devleeschouwer MJ, Barel AO. Evaluation of the preservative properties of *Thymus vulgaris* essential oil in topically applied formulations under a challenge test. *J Appl Microbiol*. 1998;84(3):368–76.
- Lambert RJ, Pearson J. Susceptibility testing: accurate and reproducible minimum inhibitory concentration (MIC) and non-inhibitory concentration (NIC) values. *J Appl Microbiol*. 2000 May;88(5):784–90.
- Rohatgi A. WebPlotDigitizer. <http://arohatgi.info/WebPlotDigitizer/app/> Accessed June 2 2014.; 2015.
- Halmi MIE, Shukor MS, Johari WLW, Shukor MY. Mathematical modelling of the degradation kinetics of *Bacillus cereus* grown on phenol. *J Environ Bioremed Toxicol*. 2014;2(1):1–5.
- Khan ZK, Katiyar R. Potent Antifungal Activity of Garlic (*Allium sativum*) against experimental murine disseminated cryptococcosis. *Pharm Biol*. 2000 Apr 1;38(2):87–100.
- Maulianawati D, Suharni S. Antibacterial activity of *Nephrolepis biserrata* extract against *Aeromonas hydrophila* and *Vibrio parahaemolyticus*. In 2022.
- Yin L, Chen J, Wang K, Geng Y, Lai W, Huang X, et al. Study the antibacterial mechanism of cinnamaldehyde against drug-resistant *Aeromonas hydrophila* in vitro. *Microb Pathog*. 2020;145.
- Majolo C, Monteiro PC, Nascimento AVPD, Chaves FCM, Gama PE, Bizzo HR, et al. Essential oils from five Brazilian piper species as antimicrobials against strains of *Aeromonas hydrophila*. *J Essent Oil-Bear Plants*. 2019;22(3):746–61.
- Pridgeon JW, Klesius PH, Mu X, Song L. An in vitro screening method to evaluate chemicals as potential chemotherapeutics to control *Aeromonas hydrophila* infection in channel catfish. *J Appl Microbiol*. 2011;111(1):114–24.
- Mahyuni S, Radzali M, Marziah M, Johari R, Mohd Aspollah S. Studies on the production of flavonoids (quercetin and hesperitin) from callus culture of *Citrus aurantifolia* (Christm and Panzer) swingle. In: *Proceedings of the Seminar Medicinal Plants: Quality Herbal Products for Healthy Living*. Forest Research Institute Malaysia (FRIM); 2000.
- Ong SL, Kiong ALP, Poosporagi R, Hussein S. Production of Flavonoid compounds in cell cultures of *Ficus deltoidea* as influenced by medium composition. *Int J Med Aromat Plants*. 2011;1(2):62–74.
- Castro SBR, Leal CAG, Freire FR, Carvalho DA, Oliveira DF, Figueiredo HCP. Antibacterial activity of plant extracts from Brazil against fish pathogenic bacteria. *Braz J Microbiol*. 2008;39(4):756–60.
- Lee S, Najiah M, Wee W. In vitro antimicrobial activities of *Colocasia esculenta* extract against *Vibrio* spp.-short communication. *Agric Slov*. 2010;7(1):5–7.
- Mahmoud AM, El-Baky RMA, Ahmed ABF, Gad GFM. Antibacterial activity of essential oils and in combination with some standard antimicrobials against different pathogens isolated from some clinical specimens. *Am J Microbiol Res*. 2016;4(1):16–25.