

JOURNAL OF BIOCHEMISTRY, MICROBIOLOGY AND BIOTECHNOLOGY



Website: http://journal.hibiscuspublisher.com/index.php/JOBIMB/index

Biodegradation of PEGs: A review

Nubli Suhaimi¹, Mohd Arif Syed¹, Nor Aripin Shamaan² and Ahmad Razi Othman^{3*}

 ¹Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, D.E, Malaysia.
 ²Faculty of Medicine and Health Sciences, Universiti Sains Islam Malaysia, 55100 USIM, Kuala Lumpur, Malaysia.
 ³Department of Chemical Engineering, Faculty of Engineering and Built Environment, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, D.E, Malaysia.

> *Corresponding author: Dr. Ahmad Razi Othman, Department of Chemical Engineering, Faculty of Engineering and Built Environment, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, D.E, Malaysia.

Email: ahmadrazi@ukm.edu.my

HISTORY

Received: 12th February 2021 Received in revised form: 14th June 2021 Accepted: 23rd July 2021

KEYWORDS

PEG PEG-degrading bacteria Toxicity PEG-degrading enzymes Bioremediation

ABSTRACT

A variety of applications involving animals and humans make use of polyethylene glycol (PEG). Because the toxicity of various PEGs is becoming increasingly apparent and has been documented several times, their removal from polluted regions is urgently required. Several studies have shown and advocated that PEG biodegradation by microorganisms may be a viable approach for remediating PEG-contaminated landfills. As a result of Malaysia's huge number of landfills, which is likely to expand in the future, it is believed that PEG contamination would rise as well, and the use of PEG-degrading microorganisms may benefit the country. This review aimed to summarize and update information on PEG-degrading microorganisms so that the information compiled can benefit local researchers in isolating more efficient PEG-degraders. The absorption of PEGS into energy under aerobic and anaerobic systems includes a variety of enzyme chains, which will present a challenge that needs to be addressed in future isolation of PEG-degraders and the elucidation of the degradation mechanisms.

INTRODUCTION

Polyethylene glycols is utilised in various areas are hydro-soluble polymers with a common HO (CH₂CH₂O)nCH₂CH₂OH, which vary in molecular weights solely from one another. These polymers in industrial goods are so widespread that they are regarded a significant contaminant along with household trash in the environment [1]. As a result, the number and quality of human sperm, genital anomalies and the incidence of breast cancer attributable to oil-based polymers have decreased. One of the chemical by-products thought to be transmitted to women feeding their boys via breast milk is dioxin, a highly carcinogenic and poisonous by-product of the process of plastics production. The combustion of plastics, in particular PVC releases dioxins and furans into the environment. Dioxin may also be absorbed in soil and water and the water is contaminated and harmful. Therefore, traditional plastic is a significant environmental issue from production through disposal. The need for biodegradable polymers continues to grow with rising awareness of waste issues and their effects on the environment [2-10].

In addition, some researchers are looking for microorganisms capable of degrading current non-biodegradable polymers such as polyvinyl chloride, polyethylene, etc. The method in which nature recycles trash or breaks organic matter into nutrients which may be utilised by another creature is biodegradation. "Degradation" implies decay, and the prefix of the organic signifies a wide variety of dead matter-eating, recycled bacteria, fungus, insects, worms and other creatures. In nature, there is no waste since everything is recycled where waste products of one organism are food for another and provide nutrients and energy when organic waste is broken down. It is a cyclical process, in which organic elements break down much more quickly than others, but ultimately everything will deteriorate [11–17].

The biodegradation of PEG by microbes has been documented and has been suggested as the future method to remediate PEG-contaminated landfills [1,18]. Since landfills in Malaysia is numerous and would continue to increase, the contamination of PEG s expected to be increased as well. Thus, bioremediation organism must be prepared. However, the usage of foreign microbes for bioremediation has potential ecological problems and is usually outcompeted by indigenous bacteria and thus is not recommended. Hence, screening of local PEGdegrading microbes must be carried out. In this thesis, the first PEG-degrading bacterium from Malaysian soils is presented.

Polyethylene Glycol

In the European pharmaceutical sector polymerization of ethylene oxide (EO, Ethylene Glycol), which is known as the macrogols, are produced as an initial material under alkaline catalysis, either water and mono-ethylene glycol or diethylene glycol. The reaction is stopped by acid neutralising the catalyst once it has achieved the appropriate molecular weight, typically controlled via viscosity measurements as in-process control. Lactic acid is usually utilised, although it may also be found in acetic acid or others. The result is a relatively basic structure of chemical: N-H where (n) the number of e-units is HO-[CH2-CH2-O] n-H [19–22].

Nomenclature of Polyethylene Glycol

Although polyethylene oxides should technically be referred to, polyethylene glycols are usually used to refer to the substantial impact of hydroxyl end groups on the chemical characteristics and the physical properties of these molecules for products with medium molecular weights of 200 to 35,000. Polyethylene oxides are referred to only products produced by polymerisation of ethylene oxides in solvents and with molecular weight up to a million. The word "PEG" in conjunction with a number value is used as an acronym for polyglycols. The number shows the mean molecular weight within the pharmaceutical business, whereas the number in the cosmetics industry shows the number (N) of EO-units in the molecule [23–25].

As ethylene oxide is 44 in weight, the PEGs are given round values of N*44 for average molecular weight values. Unfortunately, for certain PEG molecular weights, many pharmacopoeia use alternative names. **Table 1** includes the nomenclature of British Pharmacopeia II from 1993, in addition to the European, United States and Japanese monographs. The terminology is frequently still used today, even though the monograph is no longer valid. It is anticipated that a distinct and less complicated nomenclature will be created via the worldwide harmonisation of the pharmacopoeia [26].

Table 1. Nomenclature of Polyethylene Glycol (PEG) (from [27])

		Nomen-		Nomen-	Nomen-
		clature	Nomen-	clature	clature
Mean	Clariant	European Ph.	clature USP	Japanese Ph.	British Ph.
Mwt.	Polyglykol	2001	24/NF 19	XIII	II
1500	1500 S	1500	1500	-	1540
3000	3000 S or P	3000	3000	-	-
3350	3350 S or P	3350	3350	4000	4000
4000	4000 S or P	4000	4000	-	-
6000	6000 S or P	6000	6000	-	-
8000	8000 S or P	8000	8000	6000	-

Polyethylene Glycol Properties

Non-volatile fluids at room temperature are polyethylene glycols with medial molecular weight up to 400. PEG 600 at 17 to 22°C is liquid at room temperature but pasty at lower ambient temperature, whereas PEGs with a medium molecular weight of 800 to 2000 are pasty, low melting materials. The polyethylene glycols are solid over 3000 molecular weight and are not only accessible as flak but as powder. Its solubility in water is the most significant characteristic of all PEGs, which makes them excellent for usage in many applications. In any ratio of water, liquid PEGs up to PEG 600 is miscible. However, even solid grades of PEG have high water solubility. Although the molar mass decreases somewhat, even 50% (w/w) of the PEG 35,000 may be dissolved in water at normal temperature. The solution solubility and viscosity of the electrolytes, since the PEGs are noionic compounds, are not impacted [19,20,28,29].

In hard water or other aqueous solutions of different sales, PEGs are very soluble. In some PEGs, precipitates such as phenol, cresol, resorcinol, salicylic acid, tannins, iodine, tetraiodide, or mercury chloride are formed together with other compounds. Some of these reactions may be used to identify or quantify polyethylene glycols. Furthermore, liquid PEGs may be utilised to remove some of the harmful chemicals from burnt skin as antidotes. The solvent capacity of many compounds which are water-sparingly soluble is another extremely significant characteristic. The development of a compound between polyethylene glycol and the active chemical may be attributed to this activity. Polyethylene glycol has excellent acute and chronic oral toxicity, embryotoxicity or compatibility of skin. For many years they have been utilised in the cosmetics, food and pharmaceutical sector. All relevant pharmaceuticals are registered. Acceptable daily intake of polyethylene glycols (ADI) in foodstuffs with a maximum body weight of up to 10 mg/kg [21,28-30].

The Applications of Polyethylene Glycol

Polyesthers have a long history as speciality polymers used in their primary form in lubricants, anti-colding agents, ink and cosmetics as raw materials for synthesisation. They either are water-soluble or oily fluids that ultimately enter the systems of environment or wastewater. PEG is known as polyesters in large quantities and is produced in polyesters, most of which are utilised in order to synthesise nonionic surfactants as well as in various industrial products such as anti freezing agents, pharmaceutical products, cosmetics as well as lubricants as commodity chemicals. PEG is produced in copolymers in blocks and utilised as surfactants amphoteric.

Aliphatic ether bonds are very stable from a chemical point of view. In addition, the xenobiotic molecules PEG and other polyesters are genuine, given that in nature there is no tight relationship. Fortunately, no reports were discovered on polyester buildup in nature, indicating that microbes are completely metabolised [23,31]. In the pharmaceutical business, polyethylene glycol plays an essential function. The comparatively small melting point promotes a method for sintering or compression. The PEG also has a plastifying action which makes the tablet weight easy to form and may counteract capping during the compression process. In tablet coatings, solid PEGs are also often utilised. PEG is also directly tolerogenic in the transplanted organ on donor antigen [32].

Toxicity of Polyethylene Glycol

Polyethylene Glycols are neutral, molecule-soluble polymers available in a broad range of molecular weight. Many researchers have employed them as an osmotic agent for whole plants or for plant tissues, cells and organelles. The use of PEG has become a common method for reducing nutrient solutions' water potential for predefined water stress without plant consumption. Many of the researchers successfully used it to different animals where a reduction in osmotic potential without any harmful adverse consequences was ascribed to PEG responses. However, its toxicity to plants was a fairly frequent issue with PEG. Sometimes this kind of toxicity has been attributed to metal ions such as aluminium or an ionic organic molecule. While ion exchange resins, gel filtration or dialysis may be used to remove these pollutants, toxicity was not always avoided. Plant roots may not be totally impermeable for PEG, and the toxicity of the roots may be caused by intake have shown that the transport takes place without breaking down and molecular size defines the pace and location of its transport and is hypothesised that PEG impede the water route and thus promote drying. Inhibited phosphorus transport throughout the xylem root was ascribed to toxic effects of PEG [33-36]. Mexal et al. indicated a poor O2 solubility even in diluted PEG levels and a delayed transfer of O2 to root produced the major harm to plants induced by PEG [37].

Besides their toxicity on entire plants, polyethylene glycols seem to be sluggish to act parasympathomimetic substances in general. When taken intravenously, the propensity of blood to coagulate is increased and, when administered quickly, the cells may get clumped and embolization leads to death. Long contact with skins of PEG 1500 and 400 in the rabbit's skin was not harmful at doses 10 g/kg of body weight, and little, if any, of the matter was absorbed by the skin.

While early Smtyh et al. study in the 1950s [38] demonstrated skin sensitivity in a few humans and in guinea pigs tested with specific polyethylene glycols, subsequent tests revealed that materials presently manufactured are neither irritating or sensitising [39,40]. In a patient who has severe allergic reactions to several drugs on a regular basis, the cause of anaphylaxis based on an extensive diagnostic workup including skin prick tests, intradermal tests (IDT), and a double-blind oral challenge are carried out. An IDT confirmed hypersensitivity to the additive polyethylene glycol in this case, implying that the reaction was caused by an Immunoglobulin E-dependent mechanism [41].

Biodegradation

Biodegradation is a natural process in which micro-organisms are transformed into biological agents. In a basic way, it is a microorganism degradation process, a biological disruption of complex materials into a simpler inorganic component such as CO₂, H₂O, NH₃, Cl, SO₄ etc [42–46]. The method enables basic inorganic chemicals to be recycled. These atoms may be bonded in their complex structure without this process because they are non-biodegradable and unable to re-join the cycles of the ecosystems.

The actions of microorganisms may be helpful in the removal or at least the reduction of toxicity to organisms for dangerous chemicals. Soil factors such as temperature, moisture, aeration, pH and organic matter have a negative impact on microbial development and activity. microbiological degradation rate [47–53]. Compared to actinomycetes, fungus, alga, and protozoa, bacteria constitute the most prevalent soil microflora. This bacterium is found on the ground surface and multiply quickly and may be adapted to different surroundings. Therefore,

natural and manufactured compounds present in the environment are essential to decompose and convert. In degrading dangerous substances like oil spill, polyaromatic hydrocarbons, heavy metals and others, the bacteria themselves play a major role. Many research have been conducted and a few xenobiotics degrading-bacteria are identified including *Pseudomonas* sp., *Ahcromobacter* sp., *Arthrobacterium* sp., *Ahcromobacter* sp., *Arthrobacter* sp. and many more were mentioned [47–53].

Fungi are the principal microorganisms found in the forest and acid surface layers, breaking down organic materials and producing the bulk of soil and organic biomass. As a potentially feasible biodegrading agent, fungi are able to convert a broad range of dangerous substances. Inhabitants of soils include two kinds of fungus, unicellular creatures called yeasts and multicellular filament fungi (for example moulds, smuts, rods and mildews. Phanerochaete sp., Penicilium sp. and Aspergillus sp. are the common fungus that are found in the biodegradation of hazardous xenobiotics like aromatic hydrocarbon, pesticides and dyes [54-62]. Actinomycetes seem like bacteria, but is also like filamentous fungus, with their thin branch filaments. In contrast to actinomycetes, the bacteria and fungus controlled the majority of soils, however, actinomycetes may break down animal and plant-tissue resistant compounds including cellulose, chitine and phospholipids and are engaged in the production of humic substances such as green manure, compost heaps, animal dung and other plant-based materials [63-70].

Bioremediation

A big environmental issue is generated by the fast industrial advances in the globe along with the widespread usage of synthetic goods. Human neglect and lack of care are damaging to the earth's ecology. In fact, azo dyes, heavy metals, polyaromatic carbon and chemical effluents accumulate and contaminate the environment as well as xenobiotics such pesticides for agriculture, oil spill offs and industrial wastes. The remediation of this issue is part of a decision to detoxify or at least reduce toxicity for chemicals in our environment. The treatment is based on bioremediation.

Bioremediation is a biodegradation concept that has been modified and developed using technology to clean up polluted areas. It is a natural process that includes living creatures, mostly microbes, in order to reduce the danger to the environment. This may extend from the simple degradation of hazardous chemicals in natural circumstances by indigenous microbes or the isolation from non-indigenous areas, giving optimum conditions to promote the degradation of substances by microorganisms [46,71–75]. As the knowledge of the whole topic improves, the efficacy and application of bioremediation is increasing quickly. More is known about microorganisms' interacting with the physical medium nowadays and methods for GMOs are constantly developing. Constantly optimised are the techniques used to discover and isolate biodegradation enzymes.

Dealing with Plastics Waste Management

On the average, each Malaysian produces approximately 0.8 kg rubbish daily with higher figures of about 1.5 kg rubbish per day for those living in towns. According to the statistical data produced by the Ministry of Housing and Local Government in 2001, 16,247 metric ton of rubbish is produced daily. The state of Selangor leads other states in Malaysia producing about 2,995 metric ton of wastes every day. This is followed by Kuala Lumpur, which produces 2634 metric ton per day, Johor is third with an amount of 2002 metric ton per day, Perak (1596/day) and Kedah (1383/ day) [76].

From the data mentioned above, the production of wastes or solid wastes are dependent on the size of the area, the number of residents and their activities. The type of wastes being produced depends on the area itself. The solid wastes that are being produced in the town area are more complex compared from the suburban area. The products of waste are in the form of papers. glass materials, woods, plastic materials, food wastes, aluminium and so forth. Many of the trash is either transported to the waste dump site or to incinerators with 161 waste disposal settlements in Malaysia. Both systems in Malaysia are extremely popular. There is the greatest number of sites in the state of Sarawak with 28 sites. Following are Johor (26 locations), Perak (18 locations), Pahang (14 locations) and Kelantan (12 sites). Although Malaysia has a lot of garbage deposits, the growing volume of waste is still not sufficient, and the majority of sites will be overwhelmed within two years. The site will raise the danger to the environment and to the people living around the site itself. In this specific region, for example, the waste dumps may influence the land system and water [77-79].

Currently, Malaysia uses a burning technique to solve the rubbish problem caused by the limited space. Currently there are 7 mini burning centers located at four islands namely Langkawi, Tioman, Pangkor and Labuan with each of these centers has a capacity to treat about 3-10 ton/day/unit. Approximately 1,500 tons of solid wastes can be managed by this future center daily. Meanwhile, Labuan also has another thermal center which is uses the "Thermal Oxidation" method and which can handle about 40 tons of solid wastes per day. The existence of this thermal or burning center could overcome the problems of solid wastes in Malaysia [77,80].

The ability of the thermal center to burn the rubbish is faster than the open dumping system. About 99 % of the solid waste are burnt using the high temperature of between 500-600 ° Celcius. Through this method, half of the solid waste can be burnt. The ash that is being produced is approximately 99 % safe and can be buried in any landfill sites. Besides that, the thermal or burning method also destroys the microorganism inside the wastes to avoid any spread of diseases. At the same time, this system was built to take into other considerations other factors such as haze, bad smell and also the dioxin gas which will be attributed to the environment [77,80]. It is no wonder that more researchers all over the world are searching for new biodegradable polymer as well as scrambling for an answer to accelerate the biodegradation of existing recalcitrant wastes especially petroleum-based polymeric wasters more commonly known as plastics.

Microorganism and general degradation of polymer

The polymers of heterotrophic microorganisms, including bacteria and fungus, are known as potential substrate. Polymer biodegradability relies on the molecular weight, crystallinity and physical shape. An increase in molecular weight leads to a decrease in micro-organism polymer degradation. In contrast, a polymer repeating unit's monomers, dimers and oligomers are destroyed and extracted very readily. A significant decrease in solubility owing to the large molecular weight makes microbial attacks undesirable. The reason is that in bacterium, the PEG has to be absorbed through the membrane of the cell and then destroyed by cellular enzymes. It is important to note out that the deterioration of polymers may be facilitated by a competitor biological activity [81–88].

Extracellular and intracellular depolymerases are two categories of enzymes that are engaged in biological breakdown of polymers. During the depolymerization, microorganism exoenzymes break down complex polymers in order to produce short chains or smaller molecules like oligomers, dimers and monomers which are small enough to pass through semipermeable bacterial external membrane and are then used as carbon and energy sources [89]. In contrast, deterioration in inorganic species such as CO2, H2O or CH4 are known as mineralization in finishing goods. A widely known guideline is to breakdown and mineralize more closely a polymeric structure is comparable to a natural molecule. Polymers like cellulose, chitin and pullusan are bio-syntotic and may be totally and quickly biodegraded in a broad variety of natural environments via heterotrophic micro-organisms. In addition, natural circumstances include situations in which anaerobic activities are the most important breakdown of a polymer. The biodiversity and breakdown of polymer substrates may rarely approach 100% owing to the tiny part of the polymer that is integrated in the microbial biomass, humus and other natural products [81-88].

Often environmental factors influence the dominant species of bacteria and the degrading trajectories related with polymer degradation. Aerobic microorganisms with microbial biomass, CO₂, and H₂O, the end products, are mainly responsible when O₂ is present for the breakdown of complex material. Anaerobic microbe consortia are responsible, however, for the degradation of polymers in anoxic circumstances. In certain circumstances or H₂S. CO₂ and H₂O under sulfidogenic conditions the main products will be microbial biomass, CO2, CH4 and H2O. Aerobic processes generate energy that is greater than anaerobic. Aerobic energy is able to sustain a larger population of microorganisms than anaerobic ones, since thermal O2 is more efficient than SO4 and CO2 electron acceptors. These circumstances are quite common in natural settings and may be induced with suitable inoculations in the laboratory. Aerobic and strictly anaerobic micro-organisms are used for degrading polymers [3,84,86,90-93] (Fig. 1).



Fig. 1. Schematic diagram of polymer degradation under aerobic and anaerobic conditions.

Microbial degradation of PEG

Polyethers microbiological breakdown on PEG using aerobic bacteria was investigated [1]. There have been several reports of PEG aerobics using bacteria on PEG of different molecular sizes, well-described examples of which may be found in the sphingomonad genus. Fincher and Payne reported on a PEG 400-using bacteria for the first time in 1962. Diversity occurs for bacteria in regard of the maximal size of the degradable PEG molecule [18]. Different microorganisms recognised or not identified and activated sludges were absorbed into PEG up to 4,000. A limited number of species assimilated the PEG with a molecular weight of more than 4,000: *P. aeruginosa* up to 20,000; soil bacteria up to 6000; *P. stutzeri* up to 13,500 *Pseudomonas* spp. and *Sphingomonas* species down to 20,000.

of more than 6,000 has been fully digested by Sphingomonas terrrae and its associate consortia (Rhizobium, Agrobacterium Methylobacterium [94]. [95]. and spp.) Instead of lipopolysaccharides in most Gram-negative bacteria. sphenolipids are present in their outer membranes. Various lipophilic. xenobiotic. assimilating. lipophilic. αhexachlorocyclohexane, 2,4-dichloro-phenoxyacetic, dibenzo-pdioxine and diphenyl ethereal, were included in the genus: (poly)chlorofenol, polycyclic aromatic hydrocarbons [96].

Although there may be a barrier to the macromolecules for polyethers, the distinctive membrane structures of this genus may be linked with the use of membranes to take up big PEGs. This has also been supported by results of the fact that Rhodopseudomonas acidophila M402 can use its alcohol dehydrogenase to oxidise PEGs to a certain size, but cannot be grown on these compounds, which allows the dehydrogenation of high PEGs (6,000 and 20,000) that could not be used as single energy and carbon extracts of 400-1,000 or 4,000-deposit bacteria [97]. Kohlweyer et al. reported that Pseudiocardia sp. strain K1, a Gram-positive actinomycete was also identified to grow at 4,000 and 8,000 PEGs, initially being insulated as a tetrahydrofuran degrader. It is an intriguing issue whether actinomycete has the same metabolic track and mechanism of larger PEG absorption, as is discovered in sphingomonads, and the only Gram positive bacteria that can metabolize PEG [98].

Three research groups reported anaerobic assimilation of PEG. The *Alcaligenas faecalis* var. denitrifiers from the PEG bacteria were isolated from Fincher and Payne [99]. However, anaerobic growth as a denitrifier has been observed, at the cost of certain free ether glycols up to PEG 300 [100]. Dwyer and Tiedje have isolated methanogenic consortia from sewage sludge [101]. The ethylene glycol monomer and polymers of as big as PEG 20,000 were degraded by the methanogen's consortiums. Schink and Stieb [102] reported the latest group to perform the anaerobic absorption of PEG. *Pelobacter venetianus* is shown to breakdown PEG up to 20,000 in size [103]. More recently, a PEG 4000-degrading and Hexavalent Molybdenum-reducing Pseudomonas putida strain Egypt-15 has been isolated.

Growth on PEG 4,000 was optimum at concentrations from 600 to 800 mg/L and higher concentrations are inhibitory. At 800 mg/L, about 75% degradation of PEG 4,000 occurred. A lag period was observed during growth on PEG of about two days. Others have also discovered a lag period in the breakdown of PEG by bacteria, which is generally shorter (about one to two days) under aerobic settings and longer (>10 days) under anaerobic conditions [104,105]. The growth was adequately modelled using the modified Gompertz model, which give parameters such as maximum specific growth rate of 2.216 d⁻¹ and a lag period of 1.45 day. Growth on PEG was optimum at pH 7.5 and 30 °C [106].

Metabolism Of Polyethylene Glycol

In 1975 the hydrolytic breakdown of PEG 20,000 to oligomers was suggested by Haines and Alexander [22]. But early investigations of PEG aerobic breakdown have previously shown that the metabolization must be carried out via oxidation [1]. Initially, three enzymes were believed to be necessary: a dehydrogenase of alcohol and the aldehyde dehydrogenase, conversion into carboxylic acid group of the terminal alcohol groups, and an enzyme breaking ether producing glyoxylate (GOA), as illustrated in **Fig. 2** as the end product. PEG dehydrogenase (PEG-DH) and an ether bonding enzyme have been purified and described accordingly [96]. This differs greatly from the findings from Obradors and Aguilar [107], which had dehydrogenation of PEG and diglycolic acid as a single periplasmic protein (DGA). PEG-using bacteria most likely get usable energy from the oxidation stages and GOA is the basis of carbon absorption pathways. However, it has been shown that PEG-DH is unusable in bacteria that do not grow on PEG but generate dehydrogenase quinohemoprotein alcohol (QH-ADH, soluble type).

The computer modelling of a 3-D structure indicates the size of the active place is big enough to accustom a large molecule of the supports such as PEG to oxidise the alcohols and aldehydes to the corresponding acids [108]. However, since it is not yet discovered in a PEG-use organism, QH-ADH may be designed to produce alcohol dehydrogenase but probably by mistake it is called a "PEG dehydrogenase." An ether bonding enzyme seems to be dehydrogenase of β -hydroxy-acid, as detailed below. These findings may address the issue of the existence of or acquisition of PEG capability via mutations. Although the origin of the PEG metabolism machine is clear, there is still another matter of how the large molecule may reach the metabolising machine through the natural outer and inner membrane barrier. A alternative route was suggested [109] but wasn't experimentally verified, using desaturation and hydrating in the form of a hemiacetal derivative.



Fig. 2. Aerobic metabolic pathway for polyethylene glycol (PEG) (from [1])

Gram-Negative bacteria were until recently solely isolated as microorganisms for PEG use. On PEG 4000 or 8000, a grampositive actinomycete was reported to develop as tetrahydrofuran-degrader [98]. It is necessary to clarify whether the actinomycete has the same metabolic trajectory and absorption mechanism as sphingomonads with the big PEG. PEG seems to be carried straight into cytoplasmic space. PEG can control the metabalic genes for anaerobic bacteria as is proposed in the cytoplasmic space. However, not big PEGs themselves, but another PEG-transduced signal, must be regulated by metabolic genes of aerobic bacteria that seem to be barrier to large molecules, as well. The genuine regulatory components for PEGs must be found in either aerobics or anaerobics. Because derivatives such as free peg and ppg may be assimilated, but dialkyl peg/monoalkyl ppg acetate can't be used in PEG/PPG applications, at least one free alcohol group to be used for metabolism is needed. PEG/PPG [95]. As many studies bby Kawai (1987) have indicated in oxidative metabolism of PEG, the oxidised metabolites were made up of dipropylene glycol. The findings indicate that similar metabolic processes in PEG and PPG work: alcohol dehydrogenases associated with a microbial respiratory chain first attack a terminal alcohol group and then divide a terminal ether bond [110–113].

PPG must contain atactic structures since it is randomly polymerized with optically active propylene oxide. Since nearly all PPGs were added to the carbon supply, all isomers had to be digested [18]. Preliminary studies on PPG-DH showed that it exists in membrane, periplasmic and cytoplasmic fractions, even if it has not explained the relevance of these sites. PPG-dH is purified, and it should be determined that the isomers can be metabolised by a broad stereo and enantioselectivity or by various PPG-DHs.

In crude extracts of *Alcaligenes denitrificans*, the enzyme activity of poly(oxytetramethylene) glycol (PTMG) dehydrogenisation has also been discovered [114], but additional investigations have not been conducted. The terminal structure of PTMG [R-O-(CH₂)₄Oh] may be oxidised to R-O-(CH₂)₃COOH and subsequently split into R-O-CH₂COOH by β -oxidation, a structure similar to that of oxidised PEG. This may indicate that the metabolism utilises the same mechanism as PEG.

In addition, both aerobe and anaerobic bacteria have to vary in the metabolic mechanism for polymers [96]. PEG acetaldehyde lyase was the only one enzyme that might participate in PEG metabolism, as proposed by [103]. This enzyme is similar to the diol dehydratase process and divides acetaldehyde as a result, as illustrated in **Fig. 3** [115]. The cytoplasm fraction showed activity of the enzyme. These authors studied and indicated that at least an un-masked terminal hydroxyl group was required for the production of hemiacetal intermediate hydroxylation, the ether-cycled enzymes in cell free extracts from *Acetobacterium* sp. and *Pelobacter venetianus*.

This idea for the breakdown of PEG by an aerobic bacterium was originally proposed by Pearce and Heydeman [116]. Acetaldehyde is referred to by anaerobic bacteria as the ultimate result of PEG breakdown. The carbon and energy source of these creatures is most likely the acetaldehyde molecule. Dwyer and Tiedje [101] further indicated that Bacteroides and Desulfovibrios were a direct metabolite of PEG. A difference was proposed, but never tested, with the breakdown of oligomers via an unknown depolymerisation process and conversion into acetaldehyde.

PEG-dehydrogenase

Research on PEG-utilisating Sphingomonads have revealed that the PEG alcohol function is oxidised by a dye-linked, membraneclosed dehydrogenase. The enzyme possesses large suspension characteristics, however PEG dehydrogenase is named in light of its oxidation capacity (PEG-DH). PEGDHs originally were reported to be a quinoprotein (PQQ-Containing) enzyme from *Sphingomonas terrae* and *Sphingomonas macrogoltabidus* [18]. However, PEG-DH sequencing showed that the enzyme was part of the flavoprotein GMC group, and one FAD molecule was linked to the homodimeric-protein protein monomer.



Fig. 3. Anaerobic metabolism of polyethylene glycol (PEG) (from [1])

Polyethylene Glycol-Degrading Enzymes

The kinetic resolution of racemic alcohols has previously revealed an intriguing characteristic of PEG-DH [117]. Applications in bioconversions as well as in analytical chemistry of these enzymes (organisations) may be envisioned. PEG-DH participation in PEG metabolism for several additional microorganisms that use PEG was indicated [1]. A dehydrogenase, which had low activities toward PEG homologs (dimer ~PEG 600), was developed by *Alkaligenes* sp. PE18, which was isolated as an Ethylene glycol (EG) monomethyl ether-utilisating bacterium [118]. Ether-alcoholic dehydrogenase reliant on NAD generated in cell-free *Alcaligene* sp MC11 not only oxidised ether-alcohol, but also PEG homologues (n=1~3 and PEG 400) [1].

Quinohemoprotein alcohol dehydrogenase

Alcohol dehydrogenases (QH-ADHs) of quinohemoprotein may be classified into soluble (type I) and membrane-bound (type II) groups. Type II of QH-ADH can be found exclusively in Gluconobacter and Acetobacter bacteria and is not accessible information on the capacity to oxidise PEG [119]. However, certain kinds of I QH-ADH purified by other PEGuncomplicated bacteria act as alcohol dehydrogenases (ADH) for PE G: Comamonas testosteroni ADH [120] Vanillyl alcohol dehydrogenase from *Rhodopseudomonas acidophila* strain M402 [121].

C. testosteronii N-terminal portion of the enzyme is very comparable to other enzymes containing PQQ, especially methanol dehydrogenases [108] whereas C-term C does not include hemocytic. Based on this fact and on the 3-D structure of dehydrogenase methanol, modelling studies have revealed that *C. testosteroni* QH-ADH has a wide active location, explaining PEG and secondary alcohol oxidation capabilities for polyvinyl alcohol dehydrogenase [108]. In addition, three kinds of I QH-ADH identified in *Pseudomonas putida* have not been active in PEG [122]. These findings indicate that certain enzymes gained a major active site via a partial mutation to broaden their xenobiotic PEG substrate selectivity.

Dehydrogenase of liver alcohol

PEG, which may be causing mono- and dicarboxylated PEG, is known to produce a deadly condition when it's absorbed [123]. The in vivo metabolism of PEg was indicated by the presence of organic PEG acids in the blood of toxic patients and an animal model. The PEG homologs' oxidation (n=1~8) was catalysed by horse liver alcohol dehydrogenase: it is hypothesised that the syndrome may have been induced by successive oxidising of ADH and aldehyde dehydrogenase. Through different alcohol dehydrogenases, PEG may be oxidised with various routes [123] in various bacteria.

Enzyme for aldehyde oxidation

Not only primary alcohols, but also aldehydes, are the source of PEG-DH activity. The cloned PEG-DH activity towards aldehydes, however, was much weaker than towards the equivalent alcohol under optimal reaction conditions; in fact it was not significant. On the other hand, PEG, both seeming membrane-associated, was produced by separate dehydrated aldehyde dehydrogenases (AL-DH, DCIP-dependent and NAD-dependent). While the purification failed, it was found that three distinct enzymes were involved in the PEG metabolism.

Splitting enzymes Ether-bond

In general, different enzyme processes may separate the link from the ether [1,124]. However, only instances relating to the splitting of an ether bond in PEG or to its derivatives where the enzymes are separated or where such an enzyme is highly probable to result from extract activity are addressed in this study.

Dehydrogenase of PEG-carboxylic acid or diglycolic acid

Studies on PEG sphingomonad degradation showed an enzyme capable of oxidising GOA-forming PEG-carboxylic acid [1]. The DGA-DH enzyme was also known as a good substratum. DGA-DH was termed a DGA dehydrogenase. The purification of the enzyme in lauryl maltoside showed that the enzyme is obviously different from PEG-DH since it was not a substrate for either PEG or any other alcohol (including diols) [125]. Apparently, for the oxidation at uterine location in the molecule it is essential to have a terminal carboxylic acid group. The fact that GOA and GA are also substrates for DGA-DH is not necessary, however. The oxidation process may thus be comparable to that of the peroxisomal, flavoprotein glycolate (EC 1.1.3.1) and flavoprotein -hydroxide dehydragenases, by transferring hydride, by electron transfer, or by covalent catalysis in the adduct of the (carbanion) substratum and cofactor molecule [126]. In all cases, the process has to be accompanied by an H₂O molecule attack so that the rearrangement of the oxidised ether bond may be carried out under the formation of both the terminal Aldéhyde (GOA) group and the end-group Alcohol Group since no O2 use was identified for the conversion of dicarbonate PEG acid (in the PEG1). Elucidation of the mechanism and the co-factor identifiers involved must wait till significant quantities of DGA-DH are purified.

A flavoprotein DGA oxidase [127], which has no periplasmic space, was found to be present in DG A-using *Rhodococcus* sp. 432. Because ferricyanides and DCIP are significantly higher than O_2 , the enzyme may be a cytoplasmic dehydromide oxidizer of DGA on the membrane and could utilise O_2 as an in vitro electron accepter. Since both GA and GOA (but not PEG) are also used for the enzyme, glycolate oxidase may also be comparable. While structural resemblance with DGA-DH is apparently excluded by several characteristics of DGA oxidase, further research is undertaken to verify that.

In DCIP-linked dehydrogenases from a *Pseudomonas* stutzeri strain that oxidises PEG directly under GOA production, there is the same ambiguity about similarities with DGA-DH [107]. Based on PAGE and activity stained by the authors, the solubilization of a detergent was not necessary for a single enzyme oxidising PEG as well as for DGA and purifying the

enzyme. Although this suggests significant differences with the sphingomonad system, comprising two distinct enzymes (PEG-DH and DGA-DH), which require addition of solubilization detergents, no conclusions can be drawn at the moment without any additional information on the molecular characteristics of the enzyme. The initial stage of the metabolism was to produce dodecanol from *Pseudomonas* sp. strain SC25A from whole cells or cell-free extracts that may grow on PEG dodecyl ether. The split mechanism has not been investigated, nevertheless [128].

PEG acetaldehyde Lyase

PEG-degrading extracts from the anaerobic bacteria have shown diol-dehydrate and the enzyme PEG-degrading that produces acetaldehyde [115]. Unfortunately, the two enzymes seemed to be highly oxygen-sensitive, which is deemed prohibitive for purification by the authors. Since some corrinoids have been promoted by PEG degradation, they hypothesised that ether bond breaking happens in the bacterium using a PEG acetaldehyde lyase similar to diol dehydratase [102,115]. The development of acetaldehyde and PEGn-1 as products was supposed to lead to the change of position (a real mutase reaction, intramolecular rearrangement) of the terminal OH PEG group to the product, in order to create an unstable hemiacetal group.

Although the idea is validated by *Pelobacter venetianus* [102], to confirm, the enzyme needs to be purified and characterised. Furthermore, these enzymes are situated on a cytoplasm; the transportation of PEG into this compartment needs to be explained. In *P. venetianus* mentioned above two porins were discovered, however were not suitable for transporting high molecular PEG across the external cell membrane [108]. Schink discovered that phenoxyethanol is stoichiometrically metabolises by *Acetobacterium* sp. (LuPhet), while phenoxyacetic acid is not [103]. The ether connection to yielding acetaldehyde was closed as a reaction product by the cell-free extracts that clearly indicated that the chemical was destroyed by the same method as PEG anaerobic degradation.

CONCLUSION

PEG is utilised in many instances involving animal and human purposes. As the toxicity of some PEGs is becoming evident and numerously reported, its removal from contaminated areas is urgently needed. PEG biodegradation by microorganisms has been reported and proposed as a potential technique for remediating PEG-contaminated landfills. Because Malaysia has a large number of landfills and is projected to grow in the future, PEG contamination is expected to rise. Various PEG-degrading bacteria have been isolated and characterized. The route of assimilation into energy in aerobic and anaerobic system involves diverse chains of enzymes and present a challenge to those investigating the degradation of PEGs.

REFERENCES

- Kawai F. The Biochemistry of Degradation of Polyethers: Critical Reviews in Biotechnology: Vol 6, No 3. Critical Reviews in Biotechnology. 1987;6(3):273–307.
- Ando Y, Pattanawihok S, Jauzein V, Wakisaka M, Shirai Y, Nishida H. An useful biomass component for simple fabrication of the honeycomb poly(l,l-lactide) film. Nanobio and Bionano. 2014;1(1).
- Bahl S, Dolma J, Jyot Singh J, Sehgal S. Biodegradation of plastics: A state of the art review. Materials Today: Proceedings. 2021 Jan 1;39:31–4.
- 4. Bernhard M, Eubeler JP, Zok S, Knepper TP. Aerobic biodegradation of polyethylene glycols of different molecular

weights in wastewater and seawater. Water Research. 2008;42(19):4791-801.

- 5. Gopferich A. Mechanisms of polymer degradation and elimination. Handbook of Biodegradable Polymers. 1997;451–71.
- Huang SJ, Roby MS, Macri CA, Cameron JA. The effects of structure and morphology on the degradation of polymers with multiple groups. Biodegradable Polymers and Plastics. 1992;
- McCarthy SP, Ranganthan A, Ma W. Advances in properties and biodegradability of co-continuous, immicisible, biodegradable, polymer blends. Macromol Symp. 1999;144(1):63–72.
- Scott G. Photo-biodegradable plastics: Their role in the protection of the environment. Polymer Degradation and Stability. 1990;29(1):135–54.
- Tsuji H, Suzuyoshi K. Environmental degradation of biodegradable polyesters 1. Poly(\(\alpha\)-caprolactone), poly[(R)-3-hydroxybutyrate], and poly(L-lactide) films in controlled static seawater. Polymer Degradation and Stability. 2002;75(2):347–55.
- Wang Y-Z, Yang K-K, Wang X-L, Zhou Q, Zheng C-Y, Chen Z-F. Agricultural application and environmental degradation of photobiodegradable polyethylene mulching films. Journal of Polymers and the Environment. 2004;12(1):7–10.
- Breslin VT. Degradation of starch-plastic composites in a municipal solid waste landfill. Journal of Environmental Polymer Degradation. 1993;1(2):127–41.
- Habib S, Iruthayam A, Abd Shukor MY, Alias SA, Smykla J, Yasid NA. Biodeterioration of Untreated Polypropylene Microplastic Particles by Antarctic Bacteria. Polymers. 2020 Nov;12(11):2616.
- Johnson KE, Pometto III AL, Nikolov ZL. Degradation of degradable starch-polyethylene plastics in a compost environment. Applied and Environmental Microbiology. 1993;59(4):1155–61.
- Shah AA, Hasan F, Hameed A, Ahmed S. Biological degradation of plastics: A comprehensive review. Biotechnology Advances. 2008;26(3):246–65.
- Sarkar S, Basak P, Adhikari B. Biodegradation of polyethylene glycol-based polyether Urethanes. Polymer - Plastics Technology and Engineering. 2011;50(1):80–8.
- Zheng Y, Yanful EK, Bassi AS. A review of plastic waste biodegradation. Critical Reviews in Biotechnology. 2005;25(4):243–50.
- Wang Y, Fan Y, Gu J-D. Dimethyl phthalate ester degradation by two planktonic and immobilized bacterial consortia. International Biodeterioration and Biodegradation. 2004;53(2):93–101.
- Kawai F. Existence of ether bond-cleaving enzyme in a polyethylene glycol-utilizing symbiotic mixed culture. FEMS Microbiology Letters. 1985 Dec 1;30(3):273–6.
- Abdalla AL, Regitano JB, Tornisielo VL, Marchese L, Peçanha MRSR, Vitti DMSS, et al. Biodegradation of polyethylene glycol (PEG) in three tropical soils using radio labelled PEG. Animal Feed Science and Technology. 2005;122(1–2):187–93.
- Bakr MA, Ali MM, Sarkar PK. Synthesis and biodegradation of succinic acid-glycerol-polyethylene glycol-200 copolymer. Journal of Polymer Materials. 1997;14(3):251–5.
- Charoenpanich J, Tani A, Kawai F. Identification of the PEGinduced proteins by 2D-gel electrophoresis and mass spectrometry in *Sphingopyxis macrogoltabida* strain 103. Chiang Mai University Journal of Natural Sciences. 2010;9(1):111–24.
- Haines JR, Alexander M. Microbial degradation of polyethylene glycols. Journal of Applied Microbiology. 1975;29(5):621–5.
 Watson GK, Jones N. The biodegradation of polyethylene glycols
- 23. Watson GK, Jones N. The biodegradation of polyethylene glycols by sewage bacteria. Water Research. 1977;11(1):95–100.
- Tani A, Charoenpanich J, Mori T, Takeichi M, Kimbara K, Kawai F. Structure and conservation of a polyethylene glycol-degradative operon in sphingomonads. Microbiology. 2007;153(2):338–46.
- Somyoonsap P, Tani A, Charoenpanich J, Minami T, Kimbara K, Kawai F. Involvement of PEG-carboxylate dehydrogenase and glutathione S-transferase in PEG metabolism by *Sphingopyxis* macrogoltabida strain 103. Applied Microbiology and Biotechnology. 2008;81(3):473–84.
- Harris JM. Laboratory Synthesis of Polyethylene Glycol Derivatives. Journal of Macromolecular Science, Part C. 1985 Aug 1;25(3):325–73.
- Henning T. Polyethylene glycols (PEGs) and the pharmaceutical industry. SÖFW-journal. 2001;127(10):28–32.
- Abass OK, Zhuo M, Zhang K. Concomitant degradation of complex organics and metals recovery from fracking wastewater: Roles of

nano zerovalent iron initiated oxidation and adsorption. Chemical Engineering Journal. 2017;328:159–71.

- Halmi MIE, Shukor MS, Masdor NA, Shamaan NA, Shukor MY. Evaluation of several mathematical models for fitting the growth of sludge microbes on PEG 600. Journal of Environmental Microbiology and Toxicology. 2015;3(1):1–5.
- Takeuchi M, Kawai F, Shimada Y, Yokota A. Taxonomic study of polyethylene glycol-utilizing bacteria: Emended description of the genus *Sphingomonas* and new descriptions of *Sphingomonas* macrogoltabidus sp. nov., *Sphingomonas sanguis* sp. nov. and *Sphingomonas terrae* sp. nov. Systematic and Applied Microbiology. 1993;16(2):227–38.
- Traverso-Soto JM, Rojas-Ojeda P, Sanz JL, González-Mazo E, Lara-Martín PA. Anaerobic degradation of alcohol ethoxylates and polyethylene glycols in marine sediments. Science of the Total Environment. 2016;544:118–24.
- Hauet T, Goujon JM, Baumert H, Petit I, Carretier M, Eugene M, et al. Polyethylene glycol reduces the inflammatory injury due to cold ischemia/reperfusion in autotransplanted pig kidneys. Kidney Int. 2002 Aug;62(2):654–67.
- Bhuiyan TF, Ahamed KU, Nahar K, Al Mahmud J, Bhuyan MB, Anee TI, et al. Mitigation of PEG-induced drought stress in rapeseed (Brassica rapa L.) by exogenous application of osmolytes. Biocatalysis and Agricultural Biotechnology. 2019 Jul 1;20:101197.
- Chazen O, Hartung W, Neumann PM. The different effects of PEG 6000 and NaCI on leaf development are associated with differential inhibition of root water transport. Plant, Cell & Environment. 1995;18(7):727–35.
- Lawlor DW. Absorption of Polyethylene Glycols by Plants and Their Effects on Plant Growth. New Phytologist. 1970;69(2):501– 13.
- Plaut Z, Federman E. A Simple Procedure to Overcome Polyethelene Glycol Toxicity on Whole Plants. Plant Physiology. 1985 Oct 1;79(2):559–61.
- Mexal J, Fisher J, Osteryoung J, Reid CP. Oxygen Availability in Polyethylene Glycol Solutions and Its Implications in Plant-Water Relations. Plant physiology. 1975 Feb 1;55:20–4.
- Smyth HF, Carpenter CP, Weil CS. The Chronic Oral Toxicology of thePolyethylene Glycols*. Journal of the American Pharmaceutical Association (Scientific ed). 1955 Jan 1;44(1):27– 30.
- Prentice DE, Majeed SK. Oral toxicity of polyethylene glycol (PEG 200) in monkeys and rats. Toxicology Letters. 1978 Jul 1;2(2):119– 22.
- Fruijtier-Pölloth C. Safety assessment on polyethylene glycols (PEGs) and their derivatives as used in cosmetic products. Toxicology. 2005 Oct 15;214(1):1–38.
- Wylon K, Dölle S, Worm M. Polyethylene glycol as a cause of anaphylaxis. Allergy, Asthma & Clinical Immunology. 2016 Dec 13;12(1):67.
- 42. Tian F, Wang Y, Guo G, Ding K, Yang F, Wang H, et al. Enhanced azo dye biodegradation at high salinity by a halophilic bacterial consortium. Bioresource Technology. 2021;326.
- Babalola M, Ayodeji A, Bamidele O, Ajele J. Biochemical characterization of a surfactant-stable keratinase purified from *Proteus vulgaris* EMB-14 grown on low-cost feather meal. Biotechnology Letters. 2020 Dec 1;
- 44. Yu Y, Yin H, Peng H, Lu G, Dang Z. Biodegradation of decabromodiphenyl ether (BDE-209) using a novel microbial consortium GY1: Cells viability, pathway, toxicity assessment, and microbial function prediction. Science of the Total Environment. 2019;668:958–65.
- 45. Pereira E, Napp AP, Allebrandt S, Barbosa R, Reuwsaat J, Lopes W, et al. Biodegradation of aliphatic and polycyclic aromatic hydrocarbons in seawater by autochthonous microorganisms. International Biodeterioration and Biodegradation. 2019;145.
- Chaudhary DK, Kim J. New insights into bioremediation strategies for oil-contaminated soil in cold environments. International Biodeterioration and Biodegradation. 2019;142:58–72.
- Sepehr S, Shahnavaz B, Asoodeh A, Karrabi M. Biodegradation of phenol by cold-tolerant bacteria isolated from alpine soils of Binaloud Mountains in Iran. Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering. 2019;54(4):367–79.

- Bedade DK, Singhal RS. Biodegradation of acrylamide by a novel isolate, *Cupriavidus oxalaticus* ICTDB921: Identification and characterization of the acrylamidase produced. Bioresource Technology. 2018 Aug 1;261:122–32.
- Furmanczyk EM, Kaminski MA, Lipinski L, Dziembowski A, Sobczak A. *Pseudomonas laurylsulfatovorans* sp. nov., sodium dodecyl sulfate degrading bacteria, isolated from the peaty soil of a wastewater treatment plant. Syst Appl Microbiol. 2018 Jul;41(4):348–54.
- Min J, Wang J, Chen W, Hu X. Biodegradation of 2-chloro-4nitrophenol via a hydroxyquinol pathway by a Gram-negative bacterium, *Cupriavidus* sp. strain CNP-8. AMB Express. 2018 Mar 20;8(1):43.
- Singh U, Arora NK, Sachan P. Simultaneous biodegradation of phenol and cyanide present in coke-oven effluent using immobilized *Pseudomonas putida* and *Pseudomonas stutzeri*. Brazilian Journal of Microbiology. 2018 Jan 1;49(1):38–44.
- Purwanti IF, Abdullah SRS, Hamzah A, Idris M, Basri H, Mukhlisin M, et al. Biodegradation of diesel by bacteria isolated from *Scirpus mucronatus* rhizosphere in diesel-contaminated sand. Advanced Science Letters. 2015;21(2):140–3.
- Khudhair AB, Hadibarata T, Yusoff ARM, Teh ZC, Adnan LA, Kamyab H. Pyrene metabolism by new species isolated from soil *Rhizoctonia zeae* SOL3. Water, Air, and Soil Pollution. 2015;226(6).
- Sing NN, Zulkharnain A, Roslan HA, Assim Z, Husaini A. Bioremediation of PCP by *Trichoderma* and *Cunninghamella* strains isolated from sawdust. Brazilian Archives of Biology and Technology. 2014;57(6):811–20.
- Gerginova M, Manasiev J, Yemendzhiev H, Terziyska A, Peneva N, Alexieva Z. Biodegradation of phenol by Antarctic strains of *Aspergillus fumigatus*. Zeitschrift für Naturforschung C. 2014;68(9–10):384–93.
- 56. Hadibarata T, Kristanti RA. Effect of surfactants and identification of metabolites on the biodegradation of fluoranthene by basidiomycetes fungal isolate *Armillaria* sp. F022. Bioprocess and Biosystems Engineering. 2014;37(4):593–600.
- Hadibarata T, Kristanti RA. Biodegradation and metabolite transformation of pyrene by basidiomycetes fungal isolate *Armillaria* sp. F022. Bioprocess and Biosystems Engineering. 2013;36(4):461–8.
- Ahmed Abedin RM, El Hanafy AA, El-Latif SA, El-Assar SA, Fadel MS. Ligninolytic oxidative system of fungal Egyptian isolates and their applications in the decolorization of industrial dyes. Biotechnology and Biotechnological Equipment. 2013;27(6):4269–75.
- Hadibarata T, Kristanti RA, Fulazzaky MA, Nugroho AE. Characterization of pyrene biodegradation by white-rot fungus *Polyporus* sp. S133. Biotechnology and Applied Biochemistry. 2012;59(6):465-70.
- Lyra ES, Moreira KA, Porto TS, Cunha MNC da, Júnior FBP, Neto BB, et al. Decolorization of synthetic dyes by basidiomycetes isolated from woods of the Atlantic Forest (PE), Brazil. World Journal of Microbiology and Biotechnology. 2009;25(8):1499–504.
- Husaini A, Roslan HA, Hii KSY, Ang CH. Biodegradation of aliphatic hydrocarbon by indigenous fungi isolated from used motor oil contaminated sites. World Journal of Microbiology and Biotechnology. 2008;24(12):2789–97.
- Kodama T, Ding L, Yoshida M, Yajima M. Biodegradation of an striazine herbicide, simazine. Journal of Molecular Catalysis - B Enzymatic. 2001;11(4–6):1073–8.
- Castillo MA, Felis N, Aragón P, Cuesta G, Sabater C. Biodegradation of the herbicide diuron by streptomycetes isolated from soil. International Biodeterioration and Biodegradation. 2006;58(3-4):196-202.
- 64. Rehfuss M, Urban J. Rhodococcus phenolicus sp. nov., a novel bioprocessor isolated actinomycete with the ability to degrade chlorobenzene, dichlorobenzene and phenol as sole carbon sources. Systematic and Applied Microbiology. 2005;28(8):695–701.
- 65. Lee M, Kim MK, Singleton I, Goodfellow, Lee ST. Enhanced biodegradation of diesel oil by a newly identified Rhodococcus baikonurensis EN3 in the presence of mycolic acid. Journal of Applied Microbiology. 2006;100(2):325–33.

- Schrijver AD, Mot RD. Degradation of pesticides by actinomycetes. Critical Reviews in Microbiology. 1999;25(2):85– 119.
- Alvarez HM. Relationship between β-oxidation pathway and the hydrocarbon-degrading profile in actinomycetes bacteria. International Biodeterioration and Biodegradation. 2003;52(1):35– 42.
- Esposito E, Paulillo SM, Manfio GP. Biodegradation of the herbicide diuron in soil by indigenous actinomycetes. Chemosphere. 1998;37(3):541–8.
- Azmi W, Sani RK, Banerjee UC. Biodegradation of triphenylmethane dyes. Enzyme and Microbial Technology. 1998;22(3):185–91.
- Zhou W, Zimmermann W. Decolorization of industrial effluents containing reactive dyes by actinomycetes. FEMS Microbiology Letters. 1993;107(2–3):157–61.
- Yakasai HM, Rahman MF, Manogaran M, Yasid NA, Syed MA, Shamaan NA, et al. Microbiological Reduction of Molybdenum to Molybdenum Blue as a Sustainable Remediation Tool for Molybdenum: A Comprehensive Review. International Journal of Environmental Research and Public Health. 2021 Jan;18(11):5731.
- Hao L, Liu Y, Chen N, Hao X, Zhang B, Feng C. Microbial removal of vanadium (V) from groundwater by sawdust used as a sole carbon source. Science of the Total Environment. 2021;751.
- Kai EX, Johari WLW, Habib S, Yasid NA, Ahmad SA, Shukor MY. The growth of the *Rhodococcus* sp. on diesel fuel under the effect of heavy metals and different concentrations of zinc. Adv Polar Sci. 2020 May 12;132–6.
- Gan C-D, Chen T, Yang J-Y. Remediation of vanadium contaminated soil by alfalfa (Medicago sativa L.) combined with vanadium-resistant bacterial strain. Environmental Technology and Innovation. 2020;20.
- 75. Marzan LW, Sultana T, Hasan MM, Mina SA, Islam MR, Rakibuzzaman AGM, et al. Characterization of furnace oil bioremediation potential of hydrocarbonoclastic bacteria isolated from petroleum contaminated sites of the Sundarbans, Bangladesh. Journal of Genetic Engineering and Biotechnology. 2017;15(1):103–13.
- Eguchi A, Isobe T, Ramu K, Tue NM, Sudaryanto A, Devanathan G, et al. Soil contamination by brominated flame retardants in open waste dumping sites in Asian developing countries. Chemosphere. 2013 Mar;90(9):2365–71.
- Chen HL, Nath TK, Chong S, Foo V, Gibbins C, Lechner AM. The plastic waste problem in Malaysia: management, recycling and disposal of local and global plastic waste. SN Appl Sci. 2021 Mar 8;3(4):437.
- Moh YC, Abd Manaf L. Overview of household solid waste recycling policy status and challenges in Malaysia. Resources, Conservation and Recycling. 2014 Jan 1;82:50–61.
- Hassan MN, Rahman RA, Chong TL, Zakaria Z, Awang M. Waste recycling in Malaysia: problems and prospects. Waste Manag Res. 2000 Aug 1;18(4):320–8.
- Abd Kadir SAS, Yin C-Y, Rosli Sulaiman M, Chen X, El-Harbawi M. Incineration of municipal solid waste in Malaysia: Salient issues, policies and waste-to-energy initiatives. Renewable and Sustainable Energy Reviews. 2013 Aug 1;24:181–6.
- Zhao R, Guo J, Song Y, Chen Z, Lu C, Han Y, et al. Mediated electron transfer efficiencies of Se(IV) bioreduction facilitated by meso-tetrakis (4-sulfonatophenyl) porphyrin. International Biodeterioration & Biodegradation. 2020 Feb 1;147:104838.
- Shi J, Zhang B, Cheng Y, Peng K. Microbial vanadate reduction coupled to co-metabolic phenanthrene biodegradation in groundwater. Water Research. 2020;186.
- Oksińska MP, Magnucka EG, Lejcuś K, Pietr SJ. Biodegradation of the cross-linked copolymer of acrylamide and potassium acrylate by soil bacteria. Environmental Science and Pollution Research. 2016;23(6):5969–77.
- Skariyachan S, Megha M, Kini MN, Mukund KM, Rizvi A, Vasist K. Selection and screening of microbial consortia for efficient and ecofriendly degradation of plastic garbage collected from urban and rural areas of Bangalore, India. Environmental monitoring and assessment. 2015;187(1):4174.
- 85. Obruca S, Marova I, Vojtova L. Biodegradation of polyetherpolyol-based polyurethane elastomeric films: Influence of partial

replacement of polyether polyol by biopolymers of renewable origin. Environmental Technology. 2011;32(9):1043-52.

- Bao M, Chen Q, Li Y, Jiang G. Biodegradation of partially hydrolyzed polyacrylamide by bacteria isolated from production water after polymer flooding in an oil field. Journal of Hazardous Materials. 2010;184(1-3):105–10.
- Ma Y, Wang L, Shao Z. Pseudomonas, the dominant polycyclic aromatic hydrocarbon-degrading bacteria isolated from Antarctic soils and the role of large plasmids in horizontal gene transfer. Environmental Microbiology. 2006;8(3):455–65.
- Bonhomme S, Cuer A, Delort A-M, Lemaire J, Sancelme M, Scott G. Environmental biodegradation of polyethylene. Polymer Degradation and Stability. 2003;81(3):441–52.
- Bachmann BM, Seebach D. Investigation of the enzymatic cleavage of diastereomeric oligo(3-hydroxybutanoates) containing two to eight HB units. A model for the stereoselectivity of PHB depolymerase from Alcaligenes faecalis T1. Macromolecules. 1999;32(6):1777–84.
- Gupta J, Rathour R, Singh R, Thakur IS. Production and characterization of extracellular polymeric substances (EPS) generated by a carbofuran degrading strain *Cupriavidus* sp. ISTL7. Bioresource Technology. 2019;282:417–24.
- Dou W, Wei D, Li H, Li H, Rahman M, Shi J-S, et al. Purification and characterisation of a bifunctional alginate lyase from novel Isoptericola halotolerans CGMCC 5336. Carbohydrate polymers. 2013 Nov 6;98:1476–82.
- Motiwalla MJ a, Punyarthi PP b, Mehta MK b, D'Souza JS a, Kelkar-Mane V b. Studies on degradation efficiency of polycaprolactone by a naturally-occurring bacterium. Journal of Environmental Biology. 2013;34(1):43–9.
- Romen F, Reinhardt S, Jendrossek D. Thermotolerant poly(3hydroxybutyrate)-degrading bacteria from hot compost and characterization of the PHB depolymerase of Schlegellella sp. KB1a. Archives of Microbiology. 2004;182(2–3):157–64.
- 94. Takeuchi M, Kawai F, Shimada Y, Yokota A. Taxonomic Study of Polyethylene Glycol-Utilizing Bacteria: Emended Description of the Genus Sphingomonas and New Descriptions of Sphingomonas macrogoltabidus sp. nov., Sphingomonas sanguis sp. nov. and Sphingomonas terrae sp. nov. Systematic and Applied Microbiology. 1993 Jul 1;16(2):227–38.
- Kawai F, Takeuchi M. Taxonomical position of newly isolated polyethylene glycol-utilizing bacteria. Journal of Fermentation and Bioengineering. 1996 Jan 1;82(5):492–4.
- Kawai F. Sphingomonads involved in the biodegradation of xenobiotic polymers. Journal of Industrial Microbiology and Biotechnology 1999 23:4. 1999;23(4):400–7.
- Kawai F, Yamanaka H. Inducible or constitutive polyethylene glycol dehydrogenase involved in the aerobic metabolism of polyethylene glycol. Journal of Fermentation and Bioengineering. 1989 Jan 1;67(4):300-2.
- Kohlweyer U, Thiemer B, Schräder T, Andreesen JR. Tetrahydrofuran degradation by a newly isolated culture of *Pseudonocardia* sp. strain K1. FEMS Microbiology Letters. 2000;186(2):301-6.
- 99. Fincher EL, Payne WJ. Bacterial Utilization of Ether Glycols. Applied Microbiology. 1962;10(6):542.
- Grant MA, Payne WJ. Anaerobic growth of Alcaligenes faecalis var. denitrificans at the expense of ether glycols and nonionic detergents. Biotechnology and Bioengineering. 1983;25(2):627– 30.
- Dwyer DF, Tiedje JM. Degradation of ethylene glycol and polyethylene glycols by methanogenic consortia. Applied and Environmental Microbiology. 1983 Jul 1;46(1):185–90.
- 102. Schink B, Stieb M. Fermentative degradation of polyethylene glycol by a strictly anaerobic, gram-negative, nonsporeforming bacterium, Pelobacter venetianus sp. nov. Appl Environ Microbiol. 1983 Jun;45(6):1905–13.
- 103. Frings J, Schramm E, Schink B. Enzymes involved in anaerobic polyethylene glycol degradation by *Pelobacter venetianus* and *Bacteroides* strain PG1. Applied and Environmental Microbiology. 1992;58(7):2164–7.
- 104. Huang Y-L, Li Q-B, Deng X, Lu Y-H, Liao X-K, Hong M-Y, et al. Aerobic and anaerobic biodegradation of polyethylene glycols using sludge microbes. Process Biochemistry. 2005;40(1):207–11.

- 105. Traverso-Soto JM, Rojas-Ojeda P, Sanz JL, González-Mazo E, Lara-Martín PA. Anaerobic degradation of alcohol ethoxylates and polyethylene glycols in marine sediments. Science of the Total Environment. 2016;544:118–24.
- 106. AbdEl-Mongy MA, Aqlima SA, Shukor MS, Hussein S, Ling APK, Shukor MY. A PEG 4000-degrading and hexavalent molybdenumreducing *Pseudomonas putida* strain Egypt-15. Journal of the National Science Foundation of Sri Lanka. 2018 Sep 30;46(3):431– 42.
- 107. Obradors N, Aguilar J. Efficient biodegradation of high-molecularweight polyethylene glycols by pure cultures of *Pseudomonas stutzeri*. Applied and Environmental Microbiology. 1991;57(8):2383–8.
- Jongejan A, Jongejan JA, Duine JA. Homology model of the quinohaemoprotein alcohol dehydrogenase from Comamonas testosteroni. Protein Engineering, Design and Selection. 1998 Mar 1;11(3):185–98.
- 109. Thélu J, Medina L, Pelmont J. Oxidation of polyoxyethylene oligomers by an inducible enzyme from Pseudomonas P 400. FEMS Microbiology Letters. 1980 Aug 1;8(4):187–90.
- 110. Kawai F. Sphingomonads involved in the biodegradation of xenobiotic polymers. Journal of Industrial Microbiology and Biotechnology. 1999 Oct 1;23(4):400–7.
- 111. Kawai F. Breakdown of plastics and polymers by microorganisms. Advances in biochemical engineering/biotechnology. 1995;52:151–94.
- Kawai F. Microbial degradation of polyethers. Applied Microbiology and Biotechnology. 2002;58(1):30–8.
- Kawai F. The biochemistry and molecular biology of xenobiotic polymer degradation by microorganisms. Bioscience, Biotechnology and Biochemistry. 2010;74(9):1743–59.
- 114. Kawai F, Moriya F. Bacterial assimilation of polytetramethylene glycol. Journal of Fermentation and Bioengineering. 1991 Jan 1;71(1):1–5.
- 115. Schramm E, Schink B. Ether-cleaving enzyme and diol dehydratase involved in anaerobic polyethylene glycol degradation by a new *Acetobacterium* sp. Biodegradation. 1991;2(2):71–9.
- 116. Pearce BA, Heydeman MTY 1980. Metabolism of Di(ethylene glycol) [2-(2'-Hydroxyethoxy)ethanol] and Other Short Poly(ethylene glycol)s by Gram-negative Bacteria. Microbiology. 1980;118(1):21-7.
- 117. Geerlof A, Tol JBA Van, Jongejan JA, Duine JA. Enantioselective Conversions of the Racemic C3-Alcohol Synthons, Glycidol (2,3-Epoxy-1-propanol), and Solketal (2,2-Dimethyl-4-(hydroxymethyl)-l,3-dioxolane) by Quinohae-moprotein Alcohol Dehydrogenases and Bacteria Containing Such Enzymes. OUP. 2014 Jan;58(6):1028–36.
- Hino M, Murooka Y, Harada T. Enzymatic Degradation of Ether-Alcohol Compounds. Journal of Fermentation Technology. 1981;59(5):347–56.
- Ameyama M, Adachi O. [76] Alcohol dehydrogenase from acetic acid bacteria, membrane-bound. Methods in Enzymology. 1982 Jan 1;89(C):450–7.
- 120. BW G, MA van K, JA D. Quinohaemoprotein alcohol dehydrogenase apoenzyme from Pseudomonas testosteroni. The Biochemical journal. 1986;234(3):611–5.
- 121. Yamanaka K, Tsuyuki Y. A New Dye-Linked Alcohol Dehydrogenase (Vanillyl Alcohol Dehydrogenase) from Rhodopseudomonas acidophila M402 Purification, Identification of Reaction Product and Substrate Specificity. http://dx.doi.org/101080/00021369198310865937. 2014;47(10):2173-83.
- 122. Toyama H, Fujii A, Matsushita K, Shinagawa E, Ameyama M, Adachi O. Three distinct quinoprotein alcohol dehydrogenases are expressed when Pseudomonas putida is grown on different alcohols. J Bacteriol. 1995 May;177(9):2442–50.
- 123. Herold DA, Rodeheaver GT, Bellamy WT, Fitton LA, Bruns DE, Edlich RF. Toxicity of topical polyethylene glycol. Toxicology and Applied Pharmacology. 1982;65(2):329–35.
- 124. White GF, Russell NJ, Tidswell EC. Bacterial scission of ether bonds. Microbiological Reviews. 1996;60(1):216.
- 125. Enokibara S, Kawai F. Purification and characterization of an ether bond-cleaving enzyme involved in the metabolism of polyethylene glycol. Journal of Fermentation and Bioengineering. 1997 Jan 1;83(6):549–54.

- Kawai F. Microbial degradation of polyethers. Applied Microbiology and Biotechnology. 2002;58(1):30–8.
- 127. Yamanaka H, Kawai F. Purification and characterization of a glycolic acid (GA) oxidase active toward diglycolic acid (DGA) produced by DGA-utilizing Rhodococcus sp. No. 432. Journal of Fermentation and Bioengineering. 1991 Jan 1;71(2):83–8.
 128. Tidswell EC, Russell NJ, White GFY 1996. Ether-bond scission in
- Tidswell EC, Russell NJ, White GFY 1996. Ether-bond scission in the biodegradation of alcohol ethoxylate nonionic surfactants by Pseudomonas sp. strain SC25A. Microbiology. 1996;142(5):1123– 31.